

Effect of Methanol Fraction of Ethanol Extract of *Dialium Guineense* Stem Bark on Cardiovascular Disease Risk Factors in Diabetic Rats

Abu OD^{1*}, Awhin EP², and Ohikhuare F³

¹Department of Biochemistry, Faculty of Life Sciences, University of Benin, Benin City, Nigeria

²Department of Medical Biochemistry, Faculty of Basic Medical Sciences, Delta State University, Abraka, Delta State, Nigeria

³Department of Chemistry and Biochemistry, College of Letters and Science, University of Wisconsin, Milwaukee, USA

*Corresponding author: Abu OD, Faculty of Life Sciences, University of Benin, Benin City, Nigeria; E-mail: osahon.abu@uniben.edu

Received: August 03, 2023; Accepted: September 29, 2023; Published: October 12, 2023



All articles published by Gnoscience are Open Access under the Creative Commons Attribution License BY-NC-SA.

Abstract

As a disease defined by the level of hyperglycemia, diabetes mellitus (DM) causes micro- and macrovascular damage. The present study evaluated the effect of methanol fraction of ethanol extract of *Dialium guineense* (MEDG) stem bark on cardiovascular disease risk factors in diabetic rats. Male Wistar rats ($n = 25$, mean weight = 215 ± 15 g) were used. The rats were divided into five groups (5 rats per group): normal control, diabetic control, metformin, MEDG (200 mg/kg body weight, bwt) and MEDG (300 mg/kg bwt) groups. Diabetes mellitus was induced via intraperitoneal injection of streptozotocin (STZ, 50 mg/kg bwt). The diabetic rats were then treated for 21 days with metformin (50 mg/kg bwt) or MEDG (200 and 300 mg/kg bwt, respectively). The results showed that STZ-induced DM significantly increased plasma concentrations of total cholesterol (TC), triacylglycerol (TG), very low-density lipoprotein cholesterol (VLDL-C), low-density lipoprotein cholesterol (LDL-C), atherogenic index of plasma (AIP), atherogenic coefficient (AC) and cardiac risk ratio (CRR), but it significantly reduced high-density lipoprotein cholesterol (HDL-C) ($p < 0.05$). However, treatment of diabetic rats with MEDG stem bark led to significant reductions in circulating levels of lipid profile (except HDL-C, which increased) as well as AIP, AC and CRR ($p < 0.05$). These results suggest that the medicinal plant extract has the capacity to protect against cardiovascular events in diabetic rats induced by STZ.

Keywords: Atherogenic coefficient; Cholesterol; Diabetes mellitus; Lipid profile; Streptozotocin.

1. Introduction

The heart is a muscular organ which pumps blood through the blood vessels of the circulatory system [1]. Blood supplies animal body with oxygen and nutrients, while removing metabolic wastes. In humans, it is located in the middle compartment of the chest between the lungs [2]-[4]. The heart is effectively a syncytium (a meshwork of cardiac muscle cells interconnected by contiguous cytoplasmic bridges) [5]-[7]. Cardiac injury induced by drugs and other chemical

Citation: Abu OD, Awhin EP, and Ohikhuare F. Effect of methanol fraction of ethanol extract of *dialium guineense* stem bark on cardiovascular disease risk factors in diabetic rats. J Bio Med Open Access. 2023;4(1):128.

substances is becoming an increasingly important concern, since it constitutes serious risk to human health [8]. Cardiac dysfunction caused by cardiotoxic agents such as STZ may lead to heart failure, myocardial ischemia, arrhythmias, hypertension, myocarditis, pericarditis, and thromboembolism [9], [10].

Diabetes mellitus (DM) is a major cause of heart disease and stroke. Death rates for heart disease and the risk of stroke are about 2–4 times higher among adults with diabetes than among those without the disease [11]. In DM, macrovascular complications results from damage to large blood vessels and are mainly cardiovascular diseases such as premature atherosclerosis, ischemic heart disease (myocardial infarction) and stroke [12].

It has been asserted that a large population of people in developing countries use herbal medicine for some aspects of their basic health care [13], [14]. Medicinal plants have proven health benefits [15]-[19]. *Dialium guineense* is a medicinal plant used ethno-botanically to treat various ailments [20], [21]. This study investigated the effect of MEDG stem on cardiovascular disease risk factors in diabetic rats.

2. Materials and Methods

2.1 Chemicals

All chemicals, solvents and reagents used in this study were of analytical grade and they were products of Sigma-Aldrich Ltd. (USA).

2.2 Plant Extraction

Dialium guineense stem barks obtained from Auchi, Edo State, Nigeria, were authenticated at the University of Benin herbarium domiciled in the Department of Plant Biology and Biotechnology, University of Benin, Benin City, Nigeria. The prepared plant specimen was deposited in the herbarium of same department (No. UBHD330).

The plant's stem bark was washed and shade-dried for 4 weeks at room temperature, and thereafter ground into powder using a blender. A portion (500 g) of powdered plant material was steeped in 5,000 mL of absolute ethanol. The resulting extract was filtered through muslin cloth and freeze-dried with a lyophilizer. The ethanol extract was subsequently fractionated with 100% methanol [22]-[27].

2.3 Experimental Animals

Mature male Wistar albino rats (n = 25, mean weight = 215 ± 15 g) were bought from the Department of Anatomy, University of Benin, Nigeria and housed in wooden cages. They were acclimatized for two weeks before commencement of the study, and had free access to feed and water.

2.4 Experimental Design

The rats were divided into five groups (5 rats/group): normal control, diabetic control, metformin, MEDG (200 mg/kg bwt) and MEDG (300 mg/kg bwt) groups. Diabetes mellitus was induced in the rats via intraperitoneal injection of 50

mg/kg bwt STZ. The diabetic rats were then treated with either the standard antidiabetic drug metformin (50 mg/kg bwt) or MEDG (200 and 300 mg/kg bwt, respectively), for 21 days.

2.5 Blood Sample Collection and Preparation

At the end of day 21 of treatment, the rats were euthanized under mild chloroform anaesthesia after an overnight fast. Blood was drawn via cardiac puncture into heparinized sample bottles and centrifuged at 2000 rpm for 10 min to obtain plasma which was used for biochemical analyses.

2.6 Biochemical Analyses

Lipid profile and cardiovascular disease risk parameters were determined in plasma [28]-[33].

2.7 Data Analysis

Data are presented as mean \pm SEM (n = 5). Statistical analysis was performed using SPSS version 21. Statistical differences between means were compared using Duncan multiple range test. Statistical significance was assumed at $p < 0.05$.

3. Results

3.1 Effect of MEDG on Organ Weight

As shown in Fig. 1, induction of DM with STZ significantly reduced the weight of rat heart and the corresponding organ/body weight ratio ($p < 0.05$). However, treatment of the diabetic rats with MEDG markedly increased the heart weight as well as the relative organ weight ($p < 0.05$).




Fig. 1. Comparison of organ and relative organ weights.

3.2 Effect of MEDG Stem Bark on Cardiovascular Disease Risk Factors

Streptozotocin (STZ)-induced DM significantly increased plasma concentrations of TC, TG, VLDL-C, LDL-C, AIP, AC and CRR, but it significantly reduced HDL-C ($p < 0.05$). However, treatment of diabetic rats with the medicinal plant extract led to significant reductions in circulating levels of lipid profile (except HDL-C, which increased) as well as AIP, AC, and CRR ($p < 0.05$). These results are shown in Figs 2 to 5).




Fig. 2. Concentrations of TC and TG in the rats.




Fig. 3. Concentrations of LDL-C and HDL-C in the rats.




Fig. 4. Concentrations of VLDL-C and AIP in the rats.




Fig. 5. Comparison of CRR and AC of rats.

4. Discussion

People with Type-2 diabetes mellitus (T2DM) are at risk for both micro- and macrovascular complications [11]. The toxicity produced by STZ impart negatively on organs such as liver, kidneys, heart and lung. The exact molecular mechanism underlying the cytotoxic effect of STZ is not well-understood, however studies suggest that the cytotoxicity could be via production of reactive oxygen species (ROS) thus inducing oxidative stress, causing DNA damage with resultant necrosis due to the DNA methylating activity of the methyl nitroso urea moiety of the drug, release of nitric oxide (NO) which inhibits aconitase activity resulting in mitochondrial dysfunction, or via inhibition of O-linked β -N-acetylglucosaminase (O-GlcNAcase) [9].

Plants are at the center of Traditional Medicine. Their use in disease management is as old as man. Medicinal plants serve as cheap alternative to orthodox medicine since they are readily available [34]-[36]. A variety of indigenous medicinal plants can effectively ameliorates the complications of DM. One of the most significant benefits of medicinal plants is their ready availability and minimal side effects [37]-[39].

Dialium guineense (Velvet Tamarind) is a medicinal plant used in Traditional Medicine for the treatment of infections [40]. It is a tall, tropical, fruit-bearing tree, belonging to the *Leguminosae* family, and has small, typically grape-sized edible fruits with brown hard inedible shells. In Africa, it is found in dense forests along the southern border of the Sahel. The plant grows naturally in West African countries, Central African Republic, and Sudan [41]. In Nigeria, it is known by different local names: *Icheku* (Igbo), *Awin* (Yoruba), *Tsamiyarkurm* (Hausa) and *Amughen* (Bini) [42], [43].

This study investigated the effect of MEDG stem bark on cardiovascular disease risk factors in diabetic rats. The results showed that STZ-induced DM significantly increased plasma concentrations of TC, TG, VLDL-C, LDL-C, AIP, AC and CRR, but it significantly reduced HDL-C. However, treatment of diabetic rats with MEDG stem bark led to significant reductions in circulating levels of lipid profile (except HDL-C, which increased) as well as AIP, AC and CRR. These results have further strengthened the observation that medicinal plants are effective in ameliorating complications of some diseases [44]-[53].

5. Conclusion

The results of this study suggest that the medicinal plant extract has the capacity to protect against cardiovascular events in STZ-induced diabetic rats.

6. Competing Interests

The authors declare that they have no conflict of interest.

7. Acknowledgement

The authors would like to thank the laboratory staff of the Department of Biochemistry, University of Benin, Nigeria, for providing laboratory support for this work.

REFERENCES

1. Benson JM, Tibbetts BM, and Thrall KD. Uptake, tissue distribution, and fate of inhaled carbon tetrachloride: comparison of rat, mouse, and hamster. *Inhal Toxicol.* 2001;13:207-217.
2. Taber CW and Venes D. *Taber's cyclopedic medical dictionary.* Davies F.A. Co. 2009; 1018-1023p.
3. Moore KL, Dalley AF, and Agur AMR. *Clinically oriented anatomy.* Wolters Kluwer Health/Lippincott William & Wilkins. 2009; 127-173p.
4. Guyton AC and Hall JE. *Textbook of Medical Physiology* (11th ed.). Philadelphia: Elsevier Saunder. 2006; 24p.
5. MacDonald M. *Your Body: The Missing Manual.* Sebastopol, CA: Pogue Press. 2009; 40p.
6. Maton A, Jean H, Charles WM, et al. *Human Biology and Health.* Englewood Cliffs, New Jersey, USA: Prentice Hall. 1993.

7. DuBose TJ. *Fetal Sonography*, pp. 263–274; Philadelphia: WB Saunders Fennema, O. (2008). *Fennema's Food Chemistry*. CRC Press Taylor & Francis. 1996; 454–455p.
8. Dessalvi CC, Deidda M, Mele D, et al. Chemotherapy-induced cardiotoxicity: New insights into mechanisms, monitoring, and prevention. *J Cardiovascu Med*. 2018;19(7):315-323.
9. Anabel SGR, Shoshan KS, Jan DH, Alphons PLV, et al. Anomer equilibrated streptozotocin solution for the induction of experimental diabetes in mice (*Musmusculus*). *J Amer Association Lab Animal Sci*. 2010;49(1):40 – 44.
10. Colombo A, Meroni CA, Cipolla CM et al. Managing cardiotoxicity of chemotherapy. *Curr Treat Options Cardiovasc Med*. 2013;15(4):410-424.
11. Centers for Disease Control. National diabetes fact sheet: National estimates and general information on diabetes and prediabetes in the United States. 2011.
12. World Health Organization. Medicines: Safety of medicines-adverse drug reactions. 2009.
13. Luper SA. Review of plants used in the treatment of liver disease: part one. *Altern Med Rev*. 1998;3:410–421.
14. Thyagarajan SP, Jayaram S, Gopalakrishnan V, et al. Herbal medicines for liver diseases in India. *J Gastroenterol Hepatol*. 2002;17:S370–376.
15. Abu OD, Onoagbe IO, and Ekugum E. Hepatotoxicity of Graded Doses of Ethanol Extract of *Dialium guineense* Stem Bark in Wistar Rats. *J Pharm Bio-Med Sci*. 2022;2(9):347-352.
16. Abu OD, Iyare HE, and Ogboi KU. Antioxidant Property of Total Saponins and Tannins of *Dialium guineense* Stem Bark in Rats Hearts Exposed to CCl₄. *J Clinl Epidemiol Toxicol*. 2022;3(3):1-4.
17. Abu OD, Olude OM, and Obayuwana HO. Effect of methanol extract of *Citrullus lanatus* seed on hematological profile and tissue histology of normal Wistar rats. *Adv Res J Med Clin Sci*. 2021;7(7):608–615.
18. Abu OD, Imafidon KE, and Obayuwana HO. Nephrotoxic and *in vivo* antioxidant effects of *citrullus lanatus* seed extract. *Biomed J Sci Technical Res*. 2021;33(5):26281-26286.
19. Abu OD, Imafidon KE, and Obayuwana O. Effect of aqueous extract of *Anacardium occidentale* leaves on blood glucose level and lipid profile of diabetic rats. *Global Sci J*. 2020;8(10):977–987.
20. Bero J, Ganfon H, Jonville MC, et al. *In vitro* antiplasmodial activity of plants used in Benin in traditional medicine to treat malaria. *J Ethnopharmacol*. 2009;122(3):439-444.
21. Kar A. Pharmacognosy and Pharmaco-biotechnology (Revised-Expanded Second Edition). New Age International Limited Publishers, New Delhi. 2007; 332-600p.
22. Abu OD, Imafidon KE, and Iribhogbe ME. Biochemical effect of aqueous leaf extract of *Icacina trichanta* Oliv. on urea, creatinine and kidney oxidative status in CCl₄-induced Wistar rats. *Nigerian J Life Sci*. 2015;5(1):85-89.
23. Abu OD, Onoagbe IO, and Obahiagbon O. Alpha amylase and alpha glucosidase inhibitory activities of extracts of *Dialium guineense* stem bark. *Int J Clin Biol Biochem*. 2020;2(1):7-10.
24. Abu OD, Onoagbe IO, and Osemwenoyenmwun O. Alpha amylase and alpha glucosidase inhibitory activities of isolated total saponins and tannins of *Dialium guineense* stem bark. *J Cellular Mol Biol Res*. 2020;1(2):1-3.
25. Abu OD, Imafidon KE, and Iribhogbe ME. Biochemical effect of aqueous leaf extract of *Icacina trichanta* Oliv. on urea, creatinine and kidney oxidative status in CCl₄-induced Wistar rats. *Nigerian J Life Sci*. 2015;5(1):85-89.
26. Abu OD, Imafidon KE, Obayuwana HO. Phytochemical, proximate, and metal content analysis of *citrullus lanatus* (watermelon) seeds. *FUDMA J Sci*. 2017;2(2):153-156.

27. Abu OD and Onoagbe IO. Biochemical effect of aqueous extract of *Dialium Guineense* stem bark on oxidative status of normal Wistar rats. *Int J Clin Biol Biochem*. 2017;1(2):15-18.
28. Reiser R, Probstfield JL, Silvers A, et al. Plasma lipid and lipoprotein response of humans to beef fat, coconut oil and safflower oil. *Am J Clin Nutr*. 1985;42(2):190-197.
29. Tietz NW, Finley PR, and Pruden EL. *Clinical Guide to Laboratory Tests*. 2nd Edition, W.B. Saunders, Philadelphia. 1990; 304-306p.
30. Lopes-Virella MF, Stone P, Ellis S, et al. Cholesterol determination in high-density lipoproteins separated by three different methods. *Clin Chem*. 1977;23(5):882- 884.
31. Friedewald WT, Levy RI, and Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem*. 1972;18 (6):499- 502.
32. Abu OD, Eromosele AI, and Osarhenomase EG. Effect of extracts of *Dialium guineense* stem bark on lipid profile and CCl₄- induced histological changes in liver of Wistar Rats. *Int J Lipids*. 2022;1(1):22-27.
33. Frohlich J and Dobiášová M. Fractional esterification rate of cholesterol and ratio of Triglycerides to HDL-cholesterol are powerful predictors of positive findings on coronary angiography. *Clin Chem*. 2003;49 (11):1873-1880.
34. Abu OD and Ikponmwosa-Eweka O. Evaluation of the potential of total saponins and Tannins of *Dialium guineense* stem bark in the amelioration of carbon tetrachloride-induced renal oxidative stress. *SAU Sci Tech J*. 2022;7(2022;(1):42-50.
35. Abu OD, Osagie AO, and Kolawole OM. Ameliorative effect of extracts of *Dialium guineense* stem bark in CCL₄- Induced Kidney Dysfunction in Wistar Rats. *Biokemistri*. 2022e;34(2):34310-34316.
36. Abu OD, Ojo I, and Ezike TV. Methanol fraction of ethanol extract of *Dialium guineense* stem bark mitigates STZ-induced oxidative stress in Rat liver. *Biomed J Sci Techn Res*. (2023c);51(2):42594-42600.
37. Abu OD and Ikponmwosa-Eweka O. Potential of total saponins and tannins isolated from the stem bark of *Dialium guineense* in the amelioration of kidney dysfunction caused by CCl₄. *J Basic Applied Med Sci*. 2022f;2(1):1-6.
38. Abu OD and Ikponmwosa-Eweka O. Potential of extracts of *Dialium guineense* stem bark in the mitigation of carbon tetrachloride-induced renal oxidative stress. *BIU J Basic Appl Sci*. 2022g;7(1):62- 69.
39. Abu OD, Iyare HE, and Omoruyi IJ. Toxic Responses of the Blood of Rats Exposed to Aqueous Extract of *Dialium guineense* stem bark. *FUDMA J Sci*. 2022h;7(2):117-120.
40. Abu OD, Onoagbe IO, and Osemwenoyenmwun O. Alpha amylase and alpha glucosidase inhibitory activities of isolated total saponins and tannins of *Dialium guineense* stem bark. *J Cellul Mol Biol Res*. 2020;1(2):1-3.
41. Cohen G, Dembie CD, and Marcus J. Measurement of catalase activity in tissue extracts. *Analyt Biochem*. 1970;34:30-38.
42. Misra HR and Fridovich I. The role of superoxide anions in the auto oxidation of epinephrine and a single assay for superoxide dismutase. *J Biol. Chem*. 1972;247:3170-3175.
43. Rotruck JT, Pope AL, Ganther HE, et al. Selenium biochemical role as a component of glutathione peroxidase. *Science*. 1973;179:588-590

44. Ebhohon SO, Ibeh RC, Ejiofor UE, et al. Hepato- and nephro-protective effects of methanol extract of *Citrullus lanatus* rind in Wistar rats fed with used motor engine oil contaminated feed. FUDMA J Sci. 2019a;3(4):246-250.
45. Ebhohon SO, Ibeh RC, Ejiofor U, et al. Aqueous extract of *Hibiscus Sabdariffa* ameliorates cadmium-induced liver and kidney injuries in male Wistar rats. J Soc Experiment Biol Nigeria. 2019b;19(1):55-59.
46. Omoregie FO, Abu OD, and Olude OM. Ameliorative effects of aqueous leaf extract of *Annona muricata* on cyanide-induced toxicity in New Zealand Rabbits. J Bioinnovation. 2020;9(6):1532-1540.
47. Okpiabhele AO, Nwanze EAC, and Abu OD. Therapeutic potential of virgin coconut oil in ameliorating diabetes mellitus and hepatotoxicity using rattus norvegicus as case study. Asian J Biol Sci. 2018;11(3):138-144.
48. Abu OD, Onoagbe IO, and Ohikhuare F. Nephrotoxic evaluation of ethanol stem bark extract of *Dialium guineense* in normal Wistar Rats. Int J Foren Med. 2022i;4(2):19-22.
49. Abu OD, Onoagbe IO, and Ekugum E. Nephrotoxic evaluation of aqueous stem bark extract of *Dialium guineense* in normal Wistar Rats. J Pharma Bio-Med Sci. 2022j;2(9):353-357.
50. Abu OD, Umar AB, and Adekanle E. Cardiotoxic Effect of Aqueous Extract of *Dialium guineense* Stem Bark in Wistar Rats. East African Schol J Agricul Life Sci. 2022k;5(9):167-172.
51. Abu OD, Odagwe UB, and Ojo AU. Cardiotoxicity of ethanol extract of *Dialium guineense* stem bark in rats. World J Pharm Life Sci. 2022l;8(11):34-39.
52. Abu OD, Alegun O, and Ifekwe JC. Renal oxidative status in diabetic Wistar Rats administered methanol fraction of ethanol extract of *Dialium guineense*. Med Clin Case Rep J. 2023;1(1):1-13.
53. Abu OD, Awhin EP, and Ifekwe JC. Liver function status of diabetic wistar rats treated with ethanol extract of *Cucumis sativus* fruit. Biomed J Scientific Tech Rese. 2023;51(2):42440-42445.

Citation: Abu OD, Awhin EP, and Ohikhuare F. Effect of methanol fraction of ethanol extract of *dialium guineense* stem bark on cardiovascular disease risk factors in diabetic rats. J Bio Med Open Access. 2023;4(1):128.